

The Effectiveness of Perineal Massage and Kegel Exercises Against Perineal Rupture of Maternity Women at Rsi Yatofa Bodak, Central Lombok Regency

I Gusti Agung Diana¹, Sri Hardiani², Rosita Khaerina³, Endah Sulistiyani⁴
Sekolah Tinggi Ilmu Kesehatan (STIKES) Mataram

Article Info

Article history:

Accepted: May 20, 2023

Published: 2 June 2023

Keywords:

Serial Studies Knowledge retention
Hands Only CPR

Abstract

Maternal mortality and morbidity are serious health problems in developing countries. According to 2018 Riskesdas data, nationally 23.2% of deliveries experienced complications, one of which was bleeding caused by perineal rupture. Perineal rupture can be prevented by doing perineal massage and Kegel exercises. The aim of this research was to determine the effectiveness of perineal massage and Kegel exercises on perineal rupture in pregnant women at RSI Yatofa, Bodak, Central Lombok Regency. This research is pre-experimental research with a quantitative approach using an intact group comparison design. The population in this study was third trimester pregnant women who underwent pregnancy and delivery checks at RSI Yatofa, Bodak, Central Lombok Regency, totaling 44 respondents, who were divided into 2 groups, namely the perineal massage group and Kegel exercises. The sampling technique in the research was accidental sampling. Based on the results of this study, it was found that in the perineal massage group 15 people (68.2%) did not experience perineal rupture, while in the Kegel exercise group 12 people (59.1%) did not experience perineal rupture. Based on the Mann-Whitney test analysis with the help of SPSS version 26 with a total of 44 respondents, the p value was > 0.05. The conclusion shows there is no statistically significant effectiveness between the massage treatment groups perineum and gymnastics Kegel.

This is an open access article under the [Lisensi Creative Commons Atribusi-BerbagiSerupa 4.0 Internasional](#)



Corresponding Author: Ahmad I

Gusti Agung Diana

STIKES Mataram

E-mail: granddiana2016@gmail.com

1. INTRODUCTION

Maternal mortality and morbidity are currently serious health problems in developing countries. According to the 2015 World Health Organization (WHO) report, the Maternal Mortality Rate (MMR) in the world is 216 per 100,000 live births or the estimated number of maternal deaths is 303,000 deaths, with the highest number being in developing countries, namely 302,000. Several countries have quite high MMRs, such as Sub-Saharan Africa with 179,000 people, South Asia with 69,000 people and Southeast Asia with 16,000 people. The maternal mortality rate in Indonesia is 190 per 100,000 live births, Vietnam 49 per 100,000 live births, Thailand 26 per 100,000 live births, Brunei 27 per 100,000 live births, and Malaysia 29 per 100,000 live births (WHO, 2014).

Based on the 2030 Sustainable Development Goals (SDGs), there are 38 targets in the health sector that must be realized. Among other things, there is a reduction in the Maternal Mortality Rate (MMR) and Infant Mortality Rate (IMR). By 2030, the target is to reduce the MMR ratio to less than 70 per 100,000 live births (Bappenas, 2021).

From 2018 Riskesdas data, nationally 23.2% of births experienced complications. The most common type of complication is breech/transverse fetal position (3.1%), second is bleeding 2.4%. The NTB Province Health Profile notes that the number of maternal deaths in NTB Province has decreased over the last 5 years. During the 2013-2017 period, there

was a decrease in the number of maternal deaths in NTB Province by 32 people, in the same period the average decrease in the number of deaths reached 8.45% per year. In Central Lombok Regency there were 24 people. (NTB Health Profile, 2017)

The Maternal Mortality Rate (MMR) in Central Lombok Regency from 2015-2019 tends to increase from 93.4 to 123.1 per 100,000 live births. The most common causes of maternal death were hypertension in pregnancy with 14 cases (46.7%), bleeding in 8 cases (26.7%), metabolic disorders in 1 case (3.3%), infection in 1 case (3.3%) and others 6 cases (20%). One of the causes of maternal death is bleeding. The most common cause of bleeding is uterine atony or absence of uterine contractions, and the second most common cause is perineal rupture.

Factors that cause perineal rupture can occur directly or indirectly. Indirectly, pregnancy is influenced by knowledge, education, prenatal care and the mother's physical condition. Meanwhile, the time of delivery is directly influenced by parity, how to push, and the condition of the perineum. Fetal factors include the weight of the newborn and the baby's presentation (Turlina, 2015). According to Riswati (2015), spontaneous perineal rupture occurs due to tension in the vaginal area during childbirth, and can also occur due to the psychological burden of facing the birth process.

Dangers and complications the consequences of perineal rupture include bleeding and infection as well as discomfort. Bleeding from a birth canal rupture can be severe, especially in second- or third-degree birth canal ruptures or if the rupture extends to the side or up the vulva, involving the clitoris. Perineal wounds can easily become infected because their location close to the anus means they are frequently contaminated with feces. Infection can also be a reason why wounds do not heal quickly, causing scar tissue to form, causing discomfort (Turlina, 2015).

Perineal ruptureIt can be prevented in various ways, including Kegel exercises during pregnancy, correct positioning during menstruation, prenatal yoga, and perineal massage. Perineal massage is a technique for massaging the perineum during pregnancy or a few weeks before giving birth. Perineal massage can increase blood flow to the perineum and vagina so that the perineum and vagina become elastic. Perineal massage also helps relax the pelvic floor muscles so that the tissue will open without resistance during labor to facilitate the birth of the baby. Perineum perineal massage and episiotomy. Perineal massage can help soften the perineal tissue (Safrudin et al, 2009).

Perineal massage can be done at a gestational age of >34 weeks or around 1-6 weeks before giving birth (Indrayani and Tuasikal, 2020). The perineal massage technique is very safe and not dangerous so it can be done every day for 5-10 minutes, but it is not recommended to do perineal massage on mothers who have urinary tract infections or sexually transmitted infections such as herpes and fungal infections. This is supported by research by Savitri et al (2014) which stated that the incidence of perineal rupture was higher in the control group compared to the intervention group.

Data from the patient register at RSI Yatofa shows that the number of patients giving birth normally during 2020 reached 486 people. Of the total patients who experienced perineal rupture during the normal delivery process, there were 110 people or 22.6% of the total patients giving birth normally. The measure to prevent uterine rupture that is carried out in the delivery room at RSI Yatofa is to perform an episiotomy, while other preventive measures or preparations such as Kegel exercises have not been carried out.

Based on the problems above, researchers are interested in conducting research with the title "Effectiveness of perineal massage and Kegel exercises on perineal rupture of pregnant women at RSI Yatofa Bodak, Central Lombok Regency".

2. METHOD

12 | The Effectiveness of Perineal Massage and Kegel Exercises Against Perineal Rupture of Maternity Women at Rsi Yatofa Bodak, Central Lombok Regency (I Gusti Agung Diana)

This research is pre-experimental research with a quantitative approach through an intact group comparison design. The population in this study were third trimester pregnant women who underwent pregnancy and delivery checks at RSI Yatofa, Bodak, Central Lombok Regency, totaling 44 respondents, who were divided into 2 groups, namely the perineal massage group and Kegel exercises. The instrument used was a questionnaire sheet perineal massage and Kegel exercises performed by the mother and labor observation sheet. The sampling technique in the study was accidental sampling and data analysis using the nonparametric Mann-Whitney test (test of difference between two independent groups).

3. RESULTS

There were 44 respondents who participated in this research. These respondents were third trimester pregnant women who underwent pregnancy and delivery checks at RSI Yatofa, Bodak, Central Lombok Regency. This number is third trimester pregnant women who are willing to fill out the questionnaire. The normality test carried out was the Shapiro Wilk normality test because the number of samples was less than 50.

Table 1. Provides an overview of the characteristics of respondents based on age

No.	Age	Amount (n)	Percentage (%)
1	< 20 years	2	4.5%
	20s.d.35 year	34	77.3%
	> 35 years	8	18.2%
2	Education elementary school	1	2.3%
	JUNIOR HIGH SCHOOL	7	15.9%
	SENIOR HIGH SCHOOL	27	61.4%
	College	9	20.5%
3	Work Housewife	24	54.5%
	ladder Civil servant/honorary	6	13.6%
	Trader	8	18.2%
	Market Employee	6	13.6%
Total		132	100%

Source: primary data

Based on table 1 regarding the distribution of 44 respondents, the majority of respondents were in the 20-to-35-year age category, namely 34 people (77.3%), the majority of respondents had a high school education level, namely 27 people (61.4%). And most of the respondents were housewives, namely 24 respondents (54.5%).

Table 2 Distribution of respondents by treatment group

Group	Number (n)	Percentage (%)
Massage perineum	22	50%
Exercise Kegel	22	50%
Total	44	100%

Source: primary data, processed 2022

Based on table 2, it shows that the respondents were divided into 2 based on treatment groups, namely 22 respondents (50%) each.

Table 3 Frequency distribution of respondents based on the incidence of perineal rupture

Incident Rupture	Frequency (n)	Percentage (%)
Rupture	16	36.4%
Not ruptured	28	63.6%
Total	44	100%

Source: primary data

Based on table 3, it shows that the majority of respondents did not experience perineal rupture, namely 28 respondents (63.6%), and 16 respondents (36.4%) experienced rupture.

Table 4 Normality Test and Mann-Whitney Test Analysis Comparison of Perineal Massage and Kegel Exercises on Perineal Rupture at RSI Yatofa, Bodak, Central Lombok.

Tests of Normality

Intervensi	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Ruptur Pijat perineum	.430	22	.000	.590	22	.000
Senam kegel	.383	22	.000	.628	22	.000

a. Lilliefors Significance Correction

Senam kegel	22	21.50	473.00
Total	44		

Test Statistics^a

	Ruptur
Mann-Whitney U	220.000
Wilcoxon W	473.000
Z	-.620
Asymp. Sig. (2-tailed)	.536

a. Grouping Variable: Intervensi

Based on the Mann-Whitney test results in Table 4 with the help of SPSS version 26 with a total of 44 respondents with a significance level of (0.05), the value of p=0.536

($p > 0.05$) was obtained. This shows that there is no difference between perineal massage and Kegel exercises on the incidence of perineal rupture in pregnant women at RSI Yatofa, Bodak.

4. DISCUSSION

1. Perineal Massage Against Perineal Rupture

Based on table 4.6 from the cross-tabulation results, it is known that 15 of the 22 respondents in the perineal massage group (68.2%) did not experience perineal rupture and 7 people experienced perineal rupture (31.8%). This is because respondents did not regularly massage the perineum 2-3 times a week during pregnancy. Perineal massage can help stretch the inner tissues under the vagina and relax the pelvic floor muscles (Sarwono, 2008). Perineal massage performed during antenatal care starting at 34 weeks of pregnancy will reduce the possibility of perineal trauma, help reduce episiotomy and the risk of perineal laceration (Indrayani and Tuasika, 2020). Wewet (2014) proved that in the intervention group who received perineal massage, 21.4% fewer perineal ruptures occurred compared to the control group who did not receive perineal massage, namely 71.4%. Research conducted by Ugwu (2018) found that mothers who received perineal massage had an intact perineum after giving birth 59% higher than mothers who did not.

Perineal massage was performed by 29.1% so the incidence of episiotomy was significantly lower with perineal massage (Ugwu et al, 2018). Perineal massage can stimulate the connective tissue and collagen in the perineum, causing the perineum to become elastic, flexible and supple when the perineum stretches during childbirth. For this reason, it is necessary to massage the perineum to help soften the perineal tissue. This theory is supported by research conducted by Kusumawati et al (2018) which concluded that the duration of the second stage was shorter in mothers who did perineal massage compared to mothers who did not do perineal massage (Indrayani and Tuasikal, 2020).

But there are several studies that show that perineal massage is not effective in preventing perineal rupture. Research conducted by Purnami RW (2019) showed that there was no significant difference between the groups that received perineal massage and those who did not receive perineal massage. In research conducted by Lisa UF and Harmiilidyia MR (2017) it was also found that statistically perineal massage was not effective in preventing perineal rupture.

2. Kegel Exercises Against Perineal Rupture

In the Kegel exercise group, 13 respondents (59.1%) did not experience perineal rupture and 9 respondents experienced perineal rupture (36.4%). This is because the pelvic floor muscles have never been stretched before, especially in primigravida pregnant women, although there are still other things that can influence them, such as maternal, fetal and helper factors, so that motivation can be given to pregnant women to do Kegel exercises regularly. during pregnancy to prevent perineal rupture.

Research conducted by Khasanah (2014) also shows that 60% of pregnant women who do Kegel exercises do not experience perineal rupture. Research conducted by Iqmy and Minhayati (2019) on mothers giving birth found that the average perineal rupture of mothers who did Kegel exercises was 0.67 with a standard deviation of 0.617, while the average perineal rupture of mothers who did not do Kegel exercises was 1.20. with a standard deviation of 0.676.

Meanwhile, according to Kristianti's research (2015), primigravida pregnant women who did Kegel exercises (6.25%) experienced grade I perineal rupture, 37.5% experienced grade II perineal rupture and 56.25% experienced grade II perineal rupture.

III. The results of the study showed that there was no relationship between Kegel exercises in pregnant women and the degree of perineal rupture.

3. Comparison of Perineal Massage and Kegel Exercises Against Perineal Rupture

After carrying out the Mann-Whitney test for the two treatment groups, it was found that there was no statistically significant difference between the perineal massage and Kegel exercise groups, even though the percentage of the perineal massage group did not experience perineal rupture (68.2%) compared to the Kegel exercises (59.1%). Based on table 4.6, you can see the test results using the Mann-Whitney test with SPSS version 26 with a total of 44 respondents with a significance level of (0.05), obtaining a value of $p=0.536$ ($p>0.05$). This means that H_a is rejected and H_0 is accepted.

This shows that there is no difference between perineal massage and Kegel exercises on the incidence of perineal rupture in pregnant women at RSI Yatofa, Bodak. Research conducted by Rahayu et al (2015) found that perineal massage is better than Kegel exercises. The research results showed that 77.8% of the perineal massage group experienced first-degree ruptures and the rest experienced more extensive lacerations, namely second and third degrees. This study concluded that perineal massage was better at reducing perineal tearing than Kegel exercises.

A comparison between perineal massage and Kegel exercises was also carried out in research by Dewi ASK et al (2020), 20 respondents were divided into 2 groups, namely 10 respondents to the perineal massage intervention and 10 respondents to the Kegel exercise intervention. The results of the bivariate analysis showed that there was a difference in the effectiveness of perineal massage with Kegel exercises regarding the incidence of perineal rupture ($p=0.046$), it can be concluded that perineal massage is more effective in reducing the incidence of perineal rupture compared to Kegel exercises.

The incidence of perineal rupture in the mothers in the perineal massage intervention group was 40% without rupture, while in the Kegel exercise intervention group 20% did not experience rupture. The difference in the degree of rupture in primigravida mothers is partly caused by the perineum becoming more elastic, the mother feeling more comfortable and able to control herself when pushing and the mother being better able to prepare mentally for the stretch of the perineum by the baby's head (Rahayu et al, 2015).

5. CONCLUSION

The conclusion shows that there is no statistically significant effectiveness between the perineal massage and Kegel exercise treatment groups.

6. BIBLIOGRAPHY

- Amilia dkk. 2016. *Pengaruh Senam Kegel Terhadap Penurunan Gejala Inkontinensia Urin Pada Lansia Wanita di Pantai Sosial Tresna Werdha Budi Sejahtera*. Stikes Suaka Insan Banjarmasin
- Arikunto, Suharsini, 2010. *Prosedur Penelitian Suatu Pendekatan Praktek*. Jakarta: Rineke Cipta
- Astuti. 2015. *Asuhan Nifas & Menyusui*. Jakarta: Erlangga.
- Astuti L.P. dkk. 2020. *Perbedaan Efektivitas Pijat Perineum dan Supercrowning Terhadap Derajat Ruptur Perineum Pada Ibu Bersalin Primipara*. Jurnal SMART Kebidanan 7 (1): 35-40.
- Budiyarti dkk. 2019. *The Effectiveness of Kegel Exercise on The Healing Process of Episiotomy Inflammation and Proliferation Phase on Postpartum Mother*. Jurnal Keperawatan Suaka Insan Vol 4 (2).

- Depkes RI. 2010. *Profil Kesehatan Indonesia 2010*. Diakses pada tanggal 10 Oktober 2021. From <http://www.depkes.go.id>
- Dewi A.S.K. dkk. 2020. *Efektivitas Pijat Perineum dan Senam Hamil Terhadap Kejadian Ruptur Perineum*. Jurnal Penelitian Terapan Kesehatan 7 (1): 51-56.
- Farrag R.E. dkk. 2016. *Effect of Postnatal Kegel Exercise on Episiotomy Pain and Wound Healing Among Primiparous Woman*. IOSR Journal of Nursing and Health Science Volume 5 (3):24-32.
- Khasanah, Umi. 2014. *Pengaruh Senam Kegel Pada Ibu Hamil Primigravida Terhadap Kejadian Ruptur Perineum Di Poned Surakarta*. Skripsi. Universitas Sebelas Maret. Surakarta
- Kristianti, Shinta. 2015. *Hubungan Senam Kegel Pada Ibu Hamil primigravida TM III terhadap Derajat ruptur Perineum DI Wilayah Puskesmas Pembantu Bandar Kidul Kota Kediri*. Jurnal Ilmu Kesehatan 3(2): 91-98.
- Lisa U. F. dan Harmilidyya M. R. 2017. *Efektivitas Pijat Perineum Terhadap Tingkat Ruptur Perineum Pada Ibu Hamil Primigravida di Wilayah Kerja Puskesmas Alue Bilie dan Suka Mulia Kecamatan Darul Makmur Kabupaten Nagan Raya*. Journal of Healthcare Technology and Medicine 3 (1): 58-67)
- Manuaba I.B.G. 2010. *Ilmu Kebidanan, Penyakit Kandungan dan Keluarga Berencana Untuk Pendidikan Bidan*. Jakarta: EGC.
- Notoatmojo, S. 2012. *Metodologi Penelitian Kesehatan*. Jakarta: Rineka Cipta.
- Notoatmojo, Soekidjo. 2010. *Metodologi Penelitian Kesehatan*. Jakarta: PT. Rineke Cipta.
- Nursalam. 2013. *Metodologi Penelitian Ilmu Keperawatan: Pendekatan Praktis*. Edisi 3. Jakarta: Salemba Medika.
- Oxorn H dkk. 2010. *Ilmu Kebidanan, Patologi dan Fisiologi Persalinan*. Yogyakarta: Yayasan Essentia Medika (YEM).
- Profil Kesehatan Propinsi NTB Tahun 2019 Panduan Manukripsi Sekolah Tinggi Ilmu Kesehatan (STIKES) Mataram Program Studi Ilmu Kebidanan (2022).
- Prawirihardjo, Sarwono. 2011. *Ilmu Kebidanan Edisi ke 2*. Jakarta: Yayasan Bina Pustaka urnami R. W., Noviyanti R. 2019. *Efektivitas Pijat Perineum Pada Ibu Hamil Terhadap Laserasi Perineum*. Jurnal Kesehatan Madani Medika Vol 10 (2): 61-68.
- Rahayu S. dkk. 2015. *Perbedaan Hasil Masase Perineum dan Kegel Exercise Terhadap Pencegahan Robekan Perineum Pada Persalinan di Bidan*. Jurnal Riset Kesehatan 4 (2): 728-733.
- Rahmawati I, Putri, R. L. 2018. *Pengaruh Senam Kegel Terhadap Tingkat Inkontinensia Urin Pada Wanota Pasca Melahirkan*. IJOHNS Volume 3 (2): 66-71.
- Sari, E. P. Dan K. D. Rimandini. (2014). *Asuhan Kebidanan Persalinan*. Jakarta: Trans Info Media Sarniwinanti. 2018. *Pengaruh Senam Kegel Terhadap Proses Involusio Uterus Pada Ibu Post Partum*. Jurnal Kebidanan dan Keperawatan Aisyiyah Volume 14 (2): 133-139
- Siswanto, dkk. 2014. *Metodologi Penelitian Kesehatan dan Kedokteran*. Yogyakarta: Bursa Ilmu.
- Sugiyono. 2010. *Statistika Untuk Penelitian*. Bandung: Alfabeta.
- Tangko Y. dkk. 2021. *Efektivitas Pijat Perineum Terhadap Kejadian Ruptur Perineum Pada Persalinan Spontan Primigravida*. Midwifery Cara Journal 2 (4): 119-129.
- Turlina, Lilin. 2015. *Hubungan Senam Hamil Dengan Terjadinya ruptur Perineum Spontan Di BPM Wiwik Azizah Said Desa Duriwetan Kecamatan Maduran Kabupaten Lamongan*. Jurnal Ilmu Kesehatan 07 (3): 16-21
- Widianti, A, T. & Proverawati. 2010. *Senam Kesehatan*. Yogyakarta. Nuha Medika Wiknyosastro, Hanifa. 2007. *Ilmu Kebidanan*. Jakarta: Yayasan Bina Pustaka Sarwono Prawirohardjo